

RESEARCH

Open Access



# Geo-demographic and socioeconomic determinants of diagnosed hypertension among urban dwellers in Ibadan, Nigeria: a community-based study

Olalekan J. Taiwo<sup>1</sup>, Joshua O. Akinyemi<sup>2</sup>, Ayodeji Adebayo<sup>3</sup>, Oluwafemi A. Popoola<sup>3</sup>, Rufus O. Akinyemi<sup>4,5,6</sup>, Onoja M. Akpa<sup>2</sup>, Paul Olowoyo<sup>7,16</sup>, Akinkunmi P. Okekunle<sup>6</sup>, Ezinne O. Uvere<sup>5,6,9</sup>, Chukwuemeka Nwimo<sup>2,3</sup>, Omotolani Titilayo Ajala<sup>5,6</sup>, Olayinka Adebajo<sup>5,6</sup>, Adewale E. Ayodele<sup>5,6</sup>, Ayodeji Salami<sup>8</sup>, Oyedunni S. Arulogun<sup>9</sup>, Olanrewaju Olaniyan<sup>10</sup>, Richard W. Walker<sup>11</sup>, Carolyn Jenkins<sup>12</sup>, Bruce Ovbiagele<sup>13\*</sup> and Mayowa Owolabi<sup>5,6,14,15\*</sup>

## Abstract

**Background** The relationship between diagnosed high blood pressure (HBP) and proximity to health facilities and noise sources is poorly understood. We investigated the associations between the number of persons diagnosed with HBP at different distance corridors of noise-generating sources (churches, mosques, bus stops, and road networks), and blood pressure monitoring outlets (healthcare facilities and pharmaceutical shops) in Ibadan, Nigeria. In addition, we investigated the likelihood of being diagnosed with HBP using distance from noise-generating sources, distance to blood pressure monitoring outlets, socio-demographic and clinical status of the participants.

**Methods** We investigated 13,531 adults from the African Rigorous Innovative Stroke Epidemiological Surveillance (ARISES) study in Ibadan. Using a Geographic Information System (GIS), the locations of healthcare facilities, pharmaceutical shops, bus stops, churches, and mosques were buffered at 100 m intervals, and coordinates of persons diagnosed with HBP were overlaid on the buffered features. The number of persons with diagnosed HBP living at every 100 m interval was estimated. Gender, occupation, marital status, educational status, type of housing, age, and income were used as predictor variables. Analysis was conducted using Spearman rank correlation and binary logistic regression at  $p < 0.05$ .

**Results** There was a significant inverse relationship between the number of persons diagnosed with HBP and distance from pharmaceutical shops ( $r = -0.818$ ), churches ( $r = -0.818$ ), mosques ( $r = -0.893$ ) and major roads ( $r = -0.667$ ). The odds of HBP were higher among the unemployed (AOR = 1.58, 95% CI: 1.11–2.24), currently married (AOR = 1.45, CI: 1.11–1.89), and previously married (1.75, CI: 1.29–2.38). The odds of diagnosed HBP increased with educational level and age group.

\*Correspondence:  
Bruce Ovbiagele  
[Bruce.Ovbiagele@va.gov](mailto:Bruce.Ovbiagele@va.gov)  
Mayowa Owolabi  
[mayowaowolabi@yahoo.com](mailto:mayowaowolabi@yahoo.com)

Full list of author information is available at the end of the article



© The Author(s) 2024. **Open Access** This article is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License, which permits any non-commercial use, sharing, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if you modified the licensed material. You do not have permission under this licence to share adapted material derived from this article or parts of it. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by-nc-nd/4.0/>.

**Conclusion** Proximity to noise sources, being unemployed and educational level were associated with diagnosed HBP. Reduction in noise generation, transmission, and exposure could reduce the burden of hypertension in urban settings.

**Keywords** High blood pressure, Noise, Faith-based organisations, Spatial analysis, Ibadan

## Introduction

High Blood Pressure (HBP) and hypertension are often used interchangeably, but they have distinct meanings. HBP is defined as systolic blood pressure  $\geq 140$  mmHg and/or diastolic blood pressure  $\geq 90$  mm Hg due to various factors (e.g., anxiety, stress, physical activity) [1], whereas, hypertension is a chronic medical condition requiring management and treatment resulting from persistent elevation of blood pressure  $\geq 140/90$  mmHg, confirmed by multiple measurements [2]. HBP affects more than 1 billion people worldwide, the majority of whom live in low- and middle-income countries [3]. HBP and hypertension are among the most important risk factors for death and cardiovascular diseases [4]. An estimated 1.28 billion adults aged 30–79 years worldwide have hypertension, most (two-thirds) living in low- and middle-income countries [5]. This condition is a leading cause of preventable deaths, contributing to 10–20% of all deaths worldwide [6]. In LMICs, the prevalence of hypertension is notably high, with studies indicating that 26.7% of adults are affected [7].

In Europe, traffic noise is responsible for 1.7 million cases of hypertension [8]. Long exposure to noise  $> 90$  dBA has been identified as one of the drivers of HBP [9, 10, 8]. Long-term or short-term effects of exposure to environmental noise may affect the autonomic and endocrine systems resulting in changes in BP regulation that then contribute to an increased risk of cardiovascular diseases [10]. A meta-analysis of 24 studies revealed that road traffic noise is associated with an elevated risk of HBP [11]. Traffic-related noise at night causes fragmentation of sleep, the elevation of stress hormone levels, and oxidative stress which ultimately promotes the development of vascular (endothelial) dysfunction, and HBP [10].

Hypertension is a growing public health concern in Nigeria, characterized by rising prevalence rates and low awareness and treatment levels. It is a significant health concern, contributing to over 10 million deaths globally each year [12] and affects approximately 46% of adults in Africa, with studies indicating a prevalence of 32% among school teachers in Calabar [13]. Young adults are also increasingly affected, with significant risk factors including age and body mass index (BMI) [13]. Nigeria has a significant prevalence of hypertension, with approximately 25.4% of the population affected and this varies spatially ranging from 20.4% in Okpok Ikpa to 38.6% in Ikire [14]. Therefore, Nigeria's hypertension rates are among the highest in Africa, since the prevalence in

other East African countries is significantly lower, such as 9.7% in Viwandani, Kenya [14].

Previous studies on the impacts of noise on cardiovascular diseases (CVDs) have focused on industrial exposure with limited attention to noise from religious organizations [15]. In a study of 200 mosques in Riyadh, there was significantly greater noise outside, compared to inside the prayer rooms. Similarly, noise levels from places of worship were higher than those set by Jordanian limits during day and nighttime [16]. The relationship between diagnosed HBP and distance from roads and religious centres (churches and mosques), which are noise sources, has not been explored, in Nigeria, where noise from religious centres and other public facilities is unregulated.

Therefore, we investigated the associations between the number of persons diagnosed with HBP at different distance corridors of noise-generating sources (churches, mosques, bus stops, and road networks), and blood pressure monitoring outlets (healthcare facilities and pharmaceutical shops) in Ibadan, Nigeria. In addition, we investigated the likelihood of being diagnosed with HBP using distance from noise-generating sources, distance to blood pressure monitoring outlets, socio-demographic and clinical status of the population.

## Materials and methods

### Data collection

A door-to-door questionnaire survey was used to collect data between 11 October 2021 and 29 August 2022 during the baseline phase of the African Rigorous Innovative Stroke Epidemiological Surveillance (ARISES) project [17]. A very high-resolution satellite image (Google Earth) showing all the buildings in the demographic surveillance sites (DSS) was converted to PDF format and loaded to Avenza Maps software (<https://www.avenza.com/avenza-maps/>) for ease of ground navigation to respective buildings. This ensured near-total case inclusion and an accurate denominator population. Enumerators interviewed the household head or his/her proxy and collected data on demographic characteristics, stroke knowledge, stroke status, ownership of household assets and information on diagnosis of stroke, high blood pressure and other related co-morbidity using a questionnaire. In all, there were 13,531 household heads or their proxy in this urban DSS. Data was stored electronically using the REDCap mobile app [18]. Diagnosed HBP was defined as a previous diagnosis of high blood pressure

(BP  $\geq$  140/90 by a trained healthcare worker using standard protocol and/or current treatment for HBP). Furthermore, data on the locations of healthcare facilities, pharmaceutical shops, churches, mosques, and bus stops were digitized from Google Maps and validated during the data collection survey.

### Data analysis

The urban DSS data were filtered to select only those who had previously been diagnosed with HBP. The geographic coordinates of all these persons were plotted in ArcGIS software. Similarly, the locations of pharmaceutical shops, healthcare facilities, churches, mosques, and bus stops were also plotted and overlaid on the locations of the persons diagnosed with HBP to assess the relationship between distance from these healthcare facilities and noise-generating sources in ArcGIS software.

ArcGIS software was used for all the spatial analyses such as the Nearest neighbour Analysis (Rn) and Kernel Density Estimation (KDE). The Average Nearest Neighbour (Rn) analysis was used to identify the distributional pattern of persons diagnosed with HBP, healthcare facilities (pharmaceutical/patient medicine shops, and hospitals/clinics), and noise-generating sources (churches, mosques, and bus stops) [19–21]. The Rn is calculated as the observed average distance divided by the expected average distance [19]. If the average distance is less than the average for a hypothetical random distribution, the distribution of the features is considered clustered, however, if it is greater than a hypothetical random distribution, it is considered dispersed, while if it is 1, it is regarded as random [21]. The hotspot of persons diagnosed with HBP was analysed using Kernel Density Estimation (KDE) [22]. The method identified localities with higher concentrations of persons diagnosed with HBP [19].

A total of 1,510 persons previously diagnosed with HBP based on their responses to the questionnaire were selected from 13,531 respondents from the urban DSS. The coordinates of diagnosed HBP patients were overlaid on the road network, the location of churches, mosques, health facilities, and pharmaceutical shops. The number of persons diagnosed with HBP, as well as those not diagnosed with HBP at every 100 m from each healthcare facility type and noise-generating sources type were enumerated for each category using a series of non-overlapping 100 m multiple ring buffers [23].

Spearman rank correlation was used to assess the statistical relationship between distance from healthcare facilities, distance from noise-generating sources and the number of persons diagnosed with HBP. However, Analysis of Variance (ANOVA) was used to explore variations in distance from healthcare facilities and distance from noise-generating sources between those diagnosed with HBP and those not diagnosed with HBP. Chi-Square

was used to explore the association between socioeconomic factors (gender, marital status, educational level, and occupation), consumption characteristics (Tobacco, Kolanut, and alcoholic consumption) and the presence of health challenges (diabetes mellitus, blood cholesterol, obesity, and stroke) between those diagnosed with HBP and those not diagnosed with HBP.

Binary logistic regression was used to assess the association between the demographic, socioeconomic, and housing variables and the likelihood of HBP diagnosis among the entire sample of 13,531 respondents at  $p < 0.05$ . The entire sample (13,531) rather than those previously diagnosed with HBP (1,510) was used because the outcome variable in the binary logistic regression was whether a person has been diagnosed (1) or otherwise (0) with HBP. Thus, the dependent variable is a binary categorical variable indicating whether a person reported to have been diagnosed with HBP or otherwise. The regression results were presented as odds ratios (OR) together with their 95% confidence intervals (CIs). All comparisons were considered to be statistically significant at  $p < 0.05$  level.

## Results

### Distributional pattern of diagnosed HBP patients, health care facility and noise-generating sources


In all, 1510 (11.16%) of the 13,531 respondents in the urban DSS (Fig. 1) had been diagnosed with HBP. The spatial distribution analysis reveals distinct patterns in the locations of individuals diagnosed with HBP and pharmaceutical shops. The locations of individuals diagnosed with HBP and the location pharmaceutical shops are clustered in the urban DSS (Table 1) [21]. Conversely, the distribution of hospitals/Clinics, churches, and bus stops are dispersed. However, mosques are randomly distributed (Table 1).

The highest concentration of persons diagnosed with HBP are in localities such as Jayode Hospital area, Ile-Aloko Mosque area, The Bride of Christ Apostolic Church, Ayegbami Mosque, Hardex Logistic area, Olounloleru Mosque, The Hope of Heaven Christ Family Church, and Highland Specialist Hospital and Somide Creative Agency, etc. (Fig. 2).

### Proximity to healthcare facilities and pharmaceutical shops and diagnosed HBP

The average distance to healthcare facilities among persons with diagnosed HBP was significantly higher ( $214.35 \pm 91.13$  m) than for those without diagnosed HBP ( $205.44 \pm 94.27$  m,  $F_{(1,13464)} = 12.037$ ,  $p = 0.001$ ). However, there was no significant relationship between the number of persons diagnosed with HBP and distance from healthcare facilities ( $r = 0.300$ ,  $p = 0.312$ ). Thus, while the average distance travelled to hospitals varied





**Fig. 1** The selected administrative wards (Urban DSS) in Ibadan, Oyo State

**Table 1** Distributional pattern of HBP, health care facilities and noise generating sources

Variables	Number	Observed mean distance	Expected mean distance	Nearest neighbor ratio (Rn)	Z-score	P-value	Pattern
HBP Persons	1510	9.671413	25.094677	0.437726	41.799148	0.000000	Clustered
Health Care Facilities	17	270.383403	208.234018	1.298459	2.354185	0.018563	Dispersed
Pharmaceutical Shops	5	90.565038	383.964557	0.235868	3.268768	0.001080	Clustered
Churches	5	574.679866	383.964557	1.496700	2.124762	0.033606	Dispersed
Mosque	12	244.424270	247.848056	0.986186	0.091547	0.927058	Random
Bus Stops	3	718.580946	383.964557	1.871477	3.727966	0.000193	Dispersed



The average distance to churches was significantly lower among persons with diagnosed HBP ( $344.39 \pm 194.44$  m) compared to those without diagnosed HBP ( $356.93 \pm 212.36$  m;  $F_{(1,13464)} = 4.754$ ,  $p = 0.029$ ). There was

The average distance from bus terminals was significantly lower  $420.27 \pm 175.40$  m in persons with diagnosed HBP than in persons without HBP ( $438.52 \pm 173.52$ ;  $F_{(1,13464)} = 14.775$ ,  $p < 0.0001$ ). However, there was no significant relationship between the number of persons with diagnosed HBP and distance from bus terminals



( $r=-0.389$ ,  $p=0.133$ ). There was no significant difference between the distance from non-major roads among those with diagnosed HBP ( $33.00 \pm 25.42$  m) and those without HBP ( $33.82 \pm 24.83$  m;  $F_{(1,13464)}=1.481$ ,  $p=0.224$ ). The average distance from major roads among persons with diagnosed HBP ( $353.16 \pm 189.00$  m) was significantly higher than for those without diagnosed HBP ( $326.88 \pm 196.59$ ;  $F_{(1,13464)}=24.132$ ,  $p<0.0001$ ). However, there was a significant inverse relationship between distance from major roads and the number of persons with diagnosed HBP ( $r=-0.667$ ,  $p=0.006$ ).

#### Association between demographic, socioeconomic, clinical characteristics and diagnosed HBP

Persons with diagnosed HBP were significantly older (age:  $50.36 \pm 17.57$  years) than those without diagnosed HBP ( $39.23 \pm 16.10$  years;  $F_{(1,13464)}=626.147$ ,  $p<0.0001$ ). There was no significant difference in mean monthly income between persons diagnosed with HBP ( $37981.49 \pm 95593.89$  naira) and those not diagnosed with HBP ( $36233.87 \pm 86496.23$  naira;  $F_{(1,13464)}=0.533$ ,  $p=0.465$ ).

Furthermore, there was no significant difference in the proportions of males (2.7%) with diagnosed HBP, compared to females (8.5%) with diagnosed HBP ( $\chi^2=2.085$ ,  $p=0.079$ ). A greater percentage (Table 2) of married persons were diagnosed with HBP. Marital status was significantly associated with diagnosed HBP ( $\chi^2=308.067$ ,  $df=3$ ,  $p<0.05$ ). Education was also found to be significantly associated with diagnosed HBP ( $\chi^2=164.773$ ,  $p<0.05$ ), with persons who had no formal education exhibiting the highest percentage and those with primary education exhibiting the lowest percentage (Table 2). There is a significant occupational difference between those with diagnosed HBP ( $\chi^2=107.841$ ,  $p<0.05$ ) and those without HBP, with professionals having the lowest percentage of persons diagnosed with HBP (Table 2).

Regarding health-related variables, diabetes mellitus was significantly associated with diagnosed HBP ( $\chi^2=181.342$ ,  $p<0.05$ ), with individuals without diabetes mellitus being less likely to have diagnosed HBP. Those with diagnosed HBP had a significantly higher proportion of high cholesterol than those not diagnosed with HBP ( $\chi^2=38.898$ ,  $p<0.0001$ ) or obesity ( $\chi^2=20.801$ ,  $p<0.0001$ ). There was a strong association between stroke and the number of diagnosed HBP ( $\chi^2=281.253$ ,  $p<0.0001$ ), with a greater proportion of respondents who had a stroke also previously diagnosed with HBP (Table 2). Furthermore, tobacco use was significantly associated with diagnosed HBP ( $\chi^2=19.692$ ,  $p<0.05$ ). Kolanut consumption was significantly associated with diagnosed HBP ( $\chi^2=255.042$ ,  $p<0.05$ ). Alcohol consumption was significantly associated with diagnosed HBP ( $\chi^2=131.259$ ,  $p=0.0001$ ; Table 2).

#### Socioeconomic determinants of diagnosed HBP

The determinants of diagnosed HBP include occupation, marital status, education, housing type, and age. Following adjustment for demographic and socio-economic characteristics, the odds of diagnosed HBP were 58% higher among unemployed persons compared to professionals (AOR=1.58, CI: 1.11, 2.24). Similarly, the currently married (AOR=1.45, CI: 1.11, 1.89) and previously married persons (AOR=1.75, CI: 1.29, 2.38) were more likely to be diagnosed with HBP. The likelihood of HBP diagnosis was highest among those with tertiary education (AOR=1.64, CI: 1.25, 2.16) relative to those with no formal education. Furthermore, the odds of diagnosed HBP increased with the age of participants (Table 3). The Binary Logistic Regression model correctly classified 88.8% of the cases.

#### Discussion

This study sought to examine the association between the location of persons with diagnosed HBP and the location of healthcare facilities (pharmaceutical shops, hospitals/clinics) and noise-generating sources (churches, mosques and bus stops). Also, we investigated the socioeconomic predictors of diagnosed HBP among the residents. We established that persons with diagnosed HBP were clustered in space. Consequently, some local factors may be responsible for the observed distributional pattern. An overlay of the locations of churches and mosques on the Kernel Density Estimation Map shows that hotspots of persons with diagnosed HBP are within the localities where all the churches and mosques are located. Similarly, the hotspots of persons with diagnosed HBP also aligned with the major roads. Hence, there is a close association between the distribution of persons with diagnosed HBP and the distribution of churches, mosques and major road alignments. However, a relatively disproportional lower number of pharmaceutical shops are located within the hotspots of persons diagnosed with HBP.

Access to healthcare is constrained by numerous geographically varying factors, including long travel times to healthcare facilities and poor transportation infrastructure [24]. Most of the hospitals/clinics in the area are primary healthcare facilities that respond to the basic healthcare needs of the people. This perhaps accounts for why there was no significant relationship between distance from hospitals/clinics and the number of persons with diagnosed HBP. However, the distribution of pharmaceutical shops was clustered because they are often located in response to population and this could explain the negative association between distance from pharmaceutical shops and the number of persons with diagnosed HBP. In addition, proximity to pharmaceutical shops could also serve as an impetus to check one's BP. This is

**Table 2** Variations in blood pressure by socioeconomic and behavioural characteristics

Variables	Options/levels	Diagnosed HBP		Total	Chi-square	Sig
		No	Yes			
Gender	Male	3117 (23.0%)	365 (2.7%)	3473 (25.7%)	2.085	0.079
	Female	8912 (65.9%)	1148 (8.5%)	10,060 (74.3%)		
Occupation	Professional	623 (4.6%)	71 (0.5%)	694 (5.1%)	107.841	0.000
	Artisans	4491 (33.2%)	385 (2.8%)	4876 (36.0%)		
	Unemployed	629(4.6%)	138 (1.0%)	767 (5.7%)		
	Others	6275 (46.4%)	919 (6.8%)	7194 (53.2%)		
Marital Status	Single	1868 (13.8%)	81 (0.6%)	1943 (14.4%)	308.067	0.000
	Married	8310 (61.4%)	962 (7.1%)	9228 (68.5%)		
	Divorced/Separated	560 (4.1%)	116 (0.9%)	676 (5.0%)		
	Widowed	1280 (9.5%)	353 (2.6%)	1625 (12.1%)		
Education	No Formal Education	1157 (8.6%)	256 (1.9%)	1413 (10.4%)	164.773	0.000
	Primary	2225 (16.4%)	388 (2.9%)	2613 (19.3%)		
	Junior Secondary	659 (4.9%)	79 (0.6%)	736 (5.5%)		
	Senior Secondary	6771 (50.0%)	631 (4.7%)	7402 (54.7%)		
	Tertiary	1206 (8.9%)	158 (1.2%)	1364 (10.1%)		
	Others	0 (0.0%)	1 (0.0%)	1 (0.0%)		
Diabetes	No Diabetes	11,930 (88.6%)	1464 (10.9%)	13,394 (99.5%)	181.342	0.000
	Have Diabetes	28 (0.2%)	44 (0.3%)	72 (0.5%)		
Previous Stroke History	No Stroke History	11,942 (88.7%)	1478 (11.0%)	13,420 (99.7%)	135.441	0.000
	Have Stroke History	16 (0.1%)	30 (0.2%)	46 (0.3%)		
Heart Disease	No Heart Disease	11,954 (88.8%)	1507 (11.2%)	13,461 (100.0%)	0.39	0.532
	Have Heart Disease	4 (0.0%)	1 (0.0%)	5 (0.0%)		
High Cholesterol (dyslipidemia)	No Dyslipidemia	11,947 (88.7%)	1496 (11.1%)	13,443 (99.8%)	38.898	0.000
	Have Dyslipidemia	11 (0.1%)	12 (0.1%)	23 (0.2%)		
Obesity	No Obesity	11,953 (88.8%)	1502 (11.2%)	13,455 (99.9%)	20.801	0.000
	Have Obesity	5 (0.0%)	6 (0.0%)	11 (0.1%)		
Risk of Stroke	No Risk of Stroke	11,950 (88.7%)	1493 (11.1%)	13,443 (99.8%)	67.605	0.000
	Have a Risk of Stroke	8 (0.1%)	15 (0.1%)	23 (0.2%)		
Nothing	No Sickness	2645 (19.6%)	1476 (11.0%)	4121 (30.6%)	3618.961	0.000
	Yes Sickness	9313 (69.2%)	32 (0.2%)	9345 (69.4%)		
Tobacco	No Tobacco	11,597 (86.1%)	1430 (10.6%)	13,027 (96.7%)	19.692	0.000
	Yes Tobacco	361 (2.7%)	78 (0.6%)	439 (3.3%)		
Kolanut	No Kolanut	11,343 (84.2%)	1381 (10.3%)	12,724 (94.5%)	255.042	0.000
	Yes Kolanut	615 (4.6%)	127 (0.9%)	742 (5.5%)		
Alcohol	No Alcohol	13,255 (79.2%)	1337 (9.9%)	11,998 (89.1%)	131.259	0.000
	Yes Alcohol	1297 (9.6%)	171 (1.3%)	1468 (10.9%)		
Have Stroke	No Stroke	11,920 (88.5%)	1443 (10.7%)	13,363 (99.2%)	281.253	0.000
	Yes Stroke	38 (0.3%)	65 (0.5%)	103 (0.8%)		
Mild Stroke	No Mild Stroke	11,913 (88.5%)	1431 (10.6%)	13,344 (99.1%)	335.299	0.000
	Yes Mild Stroke	45 (0.3%)	77 (0.6%)	122 (0.9%)		

because most of the pharmaceutical shops provide a BP checking service for a small fee to local people.

Persons with diagnosed HBP lived much closer to churches than those without diagnosed HBP. Churches are dispersed and the percentage of persons with diagnosed HBP declined with increasing distance from churches, hence, proximity to churches is associated with diagnosed HBP. Although there is no significant difference in proximity to mosques and diagnosed HBP, perhaps because mosques are randomly distributed not necessarily in response to population distribution. Thus,

the observed association between the distribution of mosques and HBP is widespread in the study.

There was no association between distance from bus stops and the number of persons with HBP. There was no difference in distance to bus terminals between persons with HBP and those without it. All respondents are within 100 m distance from non-major roads, which may imply that the neighbourhoods are walkable. Neighbourhoods with high walkability may ameliorate the risk of hypertension and this could play a significant role in improving population health [25]. However, persons with

**Table 3** Sociodemographic and housing characteristics of respondents and the odds of diagnosed hypertension

Sociodemographic and housing variables	Categories	Diagnosed HBP: n (%)	OR (95% CI)	AOR (95% CI)
Gender	Male	365 (10.5)	1.00	
	Female	1148 (11.4)	1.08 (0.95, 1.23)	
Occupation	Professional	71 (10.2)	1.00	1.00
	Artisans	385 (7.9)	0.78 (0.59, 1.02)	1.16 (0.86, 1.57)
	Currently Unemployed	138 (18.0)	1.96 (1.43, 2.67)	1.58 (1.11, 2.24)
	Others	919 (12.8)	1.40 (1.08, 1.82)	1.30 (0.97, 1.74)
Marital Status	Never Married	81 (4.2)	1.00	1.00
	Currently Married	962 (10.4)	2.70 (2.13, 3.41)	1.45 (1.11, 1.89)
	Previously married	470 (20.4)	6.08 (4.75, 7.78)	1.75 (1.29, 2.38)
Level of Education	No Formal Education	256 (18.1)	1.00	1.00
	Primary	388 (14.9)	0.78 (0.66, 0.94)	1.35 (1.11, 1.64)
	Secondary	710 (8.7)	0.41 (0.35, 0.48)	1.31 (1.07, 1.62)
	Tertiary	159 (11.7)	0.53 (0.42, 0.66)	1.64 (1.25, 2.16)
Age (years)	< 25	69 (3.4)		1.00
	25–29	131 (6.3)	1.88 (1.39, 2.53)	1.72 (1.26, 2.34)
	30–34	119 (7.1)	2.13 (1.57, 2.90)	1.83 (1.33, 2.52)
	35–39	117 (7.7)	2.42 (1.78, 3.29)	2.01 (1.44, 2.79)
	40–44	161 (11.2)	3.62 (2.70, 4.85)	3.00 (2.18, 4.12)
	45–49	137 (12.1)	4.04 (3.00, 5.46)	3.31 (2.38, 4.60)
	50–59	293 (20.3)	7.76 (5.89, 10.21)	6.27 (4.61, 8.54)
	≥ 60	486 (22.3)	8.66 (6.66, 11.27)	6.94 (5.04, 9.57)
Income	0–50,000	1303 (11.0)	1.00	
	50,001–100,000	166 (11.9)	1.06 (0.89, 1.27)	
	> 100,001	44 (13.8)	1.24 (0.89, 1.72)	
Type of Housing Unit	Compound	31 (11.9)	1.00	
	Storey Building	528 (12.0)	0.90 (0.61, 1.35)	1.04 (0.69, 1.58)
	Bungalow	1147 (13.0)	0.95 (0.61, 1.47)	0.90 (0.58, 1.42)
	Face to face	832 (10.5)	0.78 (0.52, 1.15)	0.80 (0.53, 1.21)
	Hut	5 (50.0)	7.51 (1.99, 28.31)	8.76 (2.17, 35.37)

HBP live at a greater distance away from major roads (dual carriageway) compared with those who do have HBP and the number of persons with HBP decreases with increasing distance from major roads. There is an association between road infrastructure and HBP. Exposure to ambient  $PM_{2.5}$  concentrations from bus terminals can increase BP within a period of a few days while long-term exposure might also promote the development of chronic hypertension [26]. There is an association between the level of traffic noise and increased blood pressure among cab drivers [27].

The random and dispersed patterns of churches and mosques respectively alluded to the lack of regulations guiding the establishment and approval of locations used by faith-based organizations in the study area and this lacuna exposed the majority of the population to noise pollution from these places of worship. Although churches and mosques have different distributional patterns, a greater number of people diagnosed with HBP live in their proximity. The random distributional pattern of mosques is particularly interesting because it shows that mosques are everywhere. Bus stops are dispersed because they are located in proximity to the road

network. Therefore, both intermittent and continuous exposure to noise from religious centres and road networks can exacerbate the incidence of HBP among residents and this may worsen with the increasing density of religious centres in most African countries.

This study showed a strong, inverse relationship between distance to pharmaceutical shops, churches, mosques, road networks and the number of persons with HBP. Apart from noise from roadways, higher near-roadway particulate matter ( $PM_{2.5}$ ) and black carbon (BC) exposures have been associated with elevated BP [28]. Although, primary care clinics are frequently placed in neighbourhoods to improve access [29]. It is possible to infer that these facilities were not located in consonance with population distribution but rather based on available space. This further amplified the inadequate health infrastructural planning in most developing countries. The absence of a relationship could also be due to the combination of private and public healthcare facilities in the analysis. A similar study noted that the proximity of the home to hospitals did not correlate with primary immunization completion or BP in either a hospital-based or a community clinic [29].



Noise from haphazardly distributed faith-based centres in the unplanned metropolitan area of Ibadan could predispose residents to HBP. noise levels are seldom studied at non-workplace and non-abode sites that are visited regularly, e.g., places of worship [15]. Therefore, ambient noise levels emanating from religious activities in residential neighbourhoods are an emerging environmental problem that has attracted little attention from enforcement agencies and policymakers and components of such religious noise include, but are not limited to, mosque calls, ringing of church bells, megaphones, clapping of hands, loud prayers, chanting, singing, mobile preaching, open-air crusade, night vigils and drumming [30]. Noise measurements outside some mosques in Saudi Arabia were higher than 85 dB which is the sound level at which noise-induced hearing loss (NIHL) has been shown to occur in occupational settings [15]. Noise from religious centres has been recognized as a major source of sleep disturbances and may become unbearable with the increasing density of religious centres (churches and mosques) in the coming years [31]. Apart from sleep disorders, the literature has established the link between noise exposure and HBP [32–34].

The importance of the relationship between socioeconomic and housing variables to HBP was further established in this study. Comparatively, the likelihood of having HBP was higher among the unemployed, married, and divorced persons, while the likelihood of HBP increased with education, age group and income [14]. Current attempts at addressing the persistently high number of persons with HBP have emphasised the need to improve the lifestyle of people with limited attempts at addressing their housing and environmental conditions [35]. The findings that single individuals fare better in terms of their BP compared to their married counterparts complement and build upon earlier research and corroborate other research findings in this regard. Research has linked marital conflict to heart rate and HBP in laboratory studies [36]. Nevertheless, there are pockets of studies where never-married persons had a higher risk of HBP when compared to married persons, even after adjustment for different demographic, socioeconomic and lifestyle variables [37]. Research also suggests that occupational characteristics contribute to adverse health outcomes such as CVD [38]. Persons of high occupational status are known to have higher BP, especially men [39]. High psychological demands at the workplace, may contribute to disease risk, although, having control over one's schedule and one's work may mitigate the pathogenic effects [40]. In contrast [41], found a link between unemployment and worsening health conditions. Unemployment, increasing age, and low income have also been identified as cogent risk factors for HBP in this population and this has been reported in other studies [42].

#### Acknowledgements

We thank the staff, Field coordinators, GIS, Community managers, and Community Advisory Board/Ward Health Committees of the participating sites for their dedication and commitment.

#### Author contributions

OT, JA, BO and MOO conceptualized and designed the study. OT, JA, BO and MOO curated, analysed and interpreted the data. OT and JA wrote the first draft of the manuscript. All authors edited and approved the final draft. MOO and BO are senior authors and led the funding acquisition as overall PI and MPI respectively.

#### Funding

The National Institutes of Health grants supported the study and investigators through the ARISES (R01NS115944-01), SIREN (U54HG007479), SIBS Genomics (R01NS107900), SIBS Gen Gen (R01NS107900-02S1), H3Africa CVD Supplement (3U24HG009780-03S5), CaNVAS (1R01NS114045-01), Sub-Saharan Africa Conference on Stroke (SSACS) 1R13NS115395-01A1 and Training Africans to Lead and Execute Neurological Trials Studies (TALENTS) D43TW012030.

#### Data availability

No datasets were generated or analysed during the current study.

#### Ethics declarations

##### Ethics approval and consent to participate

Used secondary data diagnosed hypertension and the data collection tools had Informed consent which was obtained from all subjects involved in the study. Ethics approval was not required for other remotely sensed data since they are in the public domain.

##### Consent for publication

Not applicable.

##### Competing interests

The authors declare no competing interests.

##### Author details

<sup>1</sup>Department of Geography, Faculty of the Social Sciences, University of Ibadan, Ibadan, Nigeria

<sup>2</sup>Department of Epidemiology and Medical Statistics, College of Medicine, University of Ibadan, Ibadan, Nigeria

<sup>3</sup>Department of Community Medicine, College of Medicine, University of Ibadan, Ibadan, Nigeria

<sup>4</sup>Neuroscience and Ageing Research Unit, College of Medicine, IAMRAT, University of Ibadan, Ibadan, Nigeria

<sup>5</sup>Center for Genomic and Precision Medicine, College of Medicine, University of Ibadan, Ibadan, Nigeria

<sup>6</sup>Department of Medicine, College of Medicine, University of Ibadan, and University College Hospital, Ibadan, Nigeria

<sup>7</sup>Department of Medicine, Afe Babalola University, Ado Ekiti, Nigeria

<sup>8</sup>Department of Pathology, College of Medicine, University of Ibadan, Ibadan, Nigeria

<sup>9</sup>Department of Health Promotion and Education, College of Medicine, University of Ibadan, Ibadan, Nigeria

<sup>10</sup>Department of Economics, Faculty of Social Sciences, University of Ibadan, Ibadan, Nigeria

<sup>11</sup>Population Health Sciences Institute, Newcastle University, Newcastle upon Tyne, UK

<sup>12</sup>College of Nursing, Medical University of South Carolina, Charleston, USA

<sup>13</sup>Northern California Institute for Research and Education, University of California, San Francisco, USA

<sup>14</sup>Lebanese American University of Beirut Lebanon, Ibadan, Nigeria

<sup>15</sup>Blossom Specialist Medical Center, Ibadan, Nigeria

<sup>16</sup>Federal Teaching Hospital, Ido-Ekiti, Ekiti State, Nigeria

Received: 12 April 2024 / Accepted: 19 November 2024

Published online: 28 February 2025

## References

1. Wiernik E, Pannier B, Czernichow S, et al. Occupational Status Moderates the Association between current perceived stress and high blood pressure. *Hypertension*. 2013;61(3):571–7. <https://doi.org/10.1161/HYPERTENSIONAHA.111.00302>.
2. Whelton PK, Carey RM, Aronow WS, et al. NMA/PCNA Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults. *J Am Coll Cardiol*. 2018;71(19):e127–248. <https://doi.org/10.1016/j.jacc.2017.11.006>.
3. Mills KT, Bundy JD, Kelly TN, et al. Global disparities of hypertension prevalence and control. *Circulation*. 2016;134(6):441–50. <https://doi.org/10.1161/CIRCULATIONAHA.115.018912>.
4. Fisher NDL, Curfman G. Hypertension—A Public Health Challenge of Global proportions. *JAMA*. 2018;320(17):1757. <https://doi.org/10.1001/jama.2018.16760>.
5. Wang Z, Lou Y, Liu Y, et al. Methodological and reporting quality of comprehensive hypertension guidelines published between 2017 and 2022. *Chin Med J (Engl)*. 2023;136(14):1735–7. <https://doi.org/10.1097/CM9.00000000000002738>.
6. Hossain MZ. Global Burden of Hypertension and Strategic Approach to overcome it. *Bangladesh Journal of Medicine*. Published Online May. 2023;24:185. <https://doi.org/10.3329/bjrm.v34i20.66124>.
7. Hou Y, Yang S. Association of risk factors for high blood pressure across 46 low- and middle-income countries: a multi-country cross-sectional analysis. *J Glob Health*. 2024;14:04087. <https://doi.org/10.7189/jogh.14.04087>.
8. Babisch W. Cardiovascular effects of noise. *Noise Health*. 2011;13(52):201. <https://doi.org/10.4103/1463-1741.80148>.
9. Zhou F, Shrestha A, Mai S, et al. Relationship between occupational noise exposure and hypertension: a cross-sectional study in steel factories. *Am J Ind Med*. 2019;62(11):961–8. <https://doi.org/10.1002/ajim.23034>.
10. Hahad O, Kröller-Schön S, Daiber A, Münzel T. The Cardiovascular effects of noise. *Dtsch Arztebl Int*. Published Online April. 2019;5. <https://doi.org/10.3238/arztebl.2019.0245>.
11. van Kempen E, Babisch W. The quantitative relationship between road traffic noise and hypertension. *J Hypertens*. 2012;30(6):1075–86. <https://doi.org/10.1097/HJH.0b013e328352ac54>.
12. Ogungbe O, Abasilim C, Huffman MD, Ojji D. Improving hypertension control in Nigeria: early policy implications from the Hypertension treatment in Nigeria program. *Glob Health Res Policy*. 2024;9(1):26. <https://doi.org/10.1186/s41256-024-00368-9>.
13. Nwevo C, Nkang J, Nwilegbarsa S, Okonkwo U, Otu A, Ameh S. Prevalence of hypertension and Associated Factors among schoolteachers in Calabar, Nigeria: a cross-sectional study. *Niger J Clin Pract*. 2024;27(7):850–8. [https://doi.org/10.4103/njcp.njcp\\_1\\_24](https://doi.org/10.4103/njcp.njcp_1_24).
14. Okello S, Muhihi A, Mohamed SF, et al. Hypertension prevalence, awareness, treatment, and control and predicted 10-year CVD risk: a cross-sectional study of seven communities in East and West Africa (SevenCEWA). *BMC Public Health*. 2020;20(1):1706. <https://doi.org/10.1186/s12889-020-09829-5>.
15. Al Shimemeri SA, Patel CB, Abdulrahman AF. Assessment of noise levels in 200 mosques in Riyadh, Saudi Arabia. *Avicenna J Med*. 2011;01(02):35–8. <https://doi.org/10.4103/2231-0770.90914>.
16. Odat S. Noise Pollution in Irbid City — Jordan. *Fluctuation Noise Lett*. 2015;14(04):1550037. <https://doi.org/10.1142/S0219477515500376>.
17. Popoola O, Ovbiagele B, Arulogun O, et al. African rigorous innovative stroke epidemiological surveillance: protocol for a community-based Mobile-Health study. *Neuroepidemiology*. 2022;56(1):17–24. <https://doi.org/10.1159/000518885>.
18. Harris AM, Loomis J, Hopkins M, Bylund J. Assessment of Radiation Safety Knowledge among Urology residents in the United States. *J Endourol*. 2019;33(6):492–7. <https://doi.org/10.1089/end.2019.0133>.
19. Carrero-Pazos M. Density, intensity and clustering patterns in the spatial distribution of galician megaliths (NW Iberian Peninsula). *Archaeol Anthropol Sci*. 2019;11(5):2097–108. <https://doi.org/10.1007/s12520-018-0662-2>.
20. Dhanaraj K, Angadi DP. A GIS based interpretation of the historical evolution of urban settlements in Mangalore City, India. *Spat Inform Res*. 2021;29(4):615–29. <https://doi.org/10.1007/s41324-020-00363-5>.
21. MANSOUR S. Spatial analysis of public health facilities in Riyadh Governorate, Saudi Arabia: a GIS-based study to assess geographic variations of service provision and accessibility. *Geo-spatial Inform Sci*. 2016;19(1):26–38. <https://doi.org/10.1080/10095020.2016.1151205>.
22. King TL, Bentley RJ, Thornton LE, Kavanagh AM. Using kernel density estimation to understand the influence of neighbourhood destinations on BMI. *BMJ Open*. 2016;6(2):e008878. <https://doi.org/10.1136/bmjopen-2015-008878>.
23. Hazrin H, Fadhli Y, Tahir A, Safurah J, Kamaliah MN, Noraini MY. Spatial patterns of health clinic in Malaysia. *Health N Hav*. 2013;05(12):2104–9. <https://doi.org/10.4236/health.2013.512287>.
24. Weiss DJ, Nelson A, Vargas-Ruiz CA, et al. Global maps of travel time to health-care facilities. *Nat Med*. 2020;26(12):1835–8. <https://doi.org/10.1038/s41591-020-1059-1>.
25. Li F, Harmer P, Cardinal BJ, Vongjaturapat N. Built environment and changes in blood pressure in middle aged and older adults. *Prev Med (Baltim)*. 2009;48(3):237–41. <https://doi.org/10.1016/j.ypmed.2009.01.005>.
26. Brook RD, Rajagopalan S. Particulate matter, air pollution, and blood pressure. *J Am Soc Hypertens*. 2009;3(5):332–50. <https://doi.org/10.1016/j.JASH.2009.08.005>.
27. Addina S, Soedjajadi K. Relationship of traffic noise with high blood pressure to Pedicab drivers around Purabaya Bus Station Surabaya. *Jurnal Kesehatan Lingkungan Unair*. 2015;8(1):69–80.
28. Morishita M, Wang L, Speth K, et al. Acute blood pressure and Cardiovascular effects of Near-Roadway exposures with and without N95 respirators. *Am J Hypertens*. 2019;32(11):1054–65. <https://doi.org/10.1093/ajh/hpz113>.
29. Baumgardner DJ, Halsmer SE, Steber DL, Shah DS, Mundt MP. Does Proximity to Clinic affect immunization rates and blood pressure? *Int J Psychiatry Med*. 2006;36(2):199–209. <https://doi.org/10.2190/9N36-W446-194L-9KHW>.
30. Armah FA, Odoi JO, Yawson DO, Yengoh GT, Afrifa EKA, Pappoe ANM. Mapping of noise risk zones derived from religious activities and perceptions in residential neighbourhoods in the Cape Coast metropolis, Ghana. *Environ Hazards*. 2010;9(4):358–68. <https://doi.org/10.3763/ehaz.2010.0003>.
31. Zakpala RN, Armah FA, Sackey BM, Pabi O. Night-time decibel hell: mapping noise exposure zones and individual annoyance ratings in an urban environment in Ghana. *Scientifica (Cairo)*. 2014;2014:1–11. <https://doi.org/10.1155/2014/892105>.
32. Gesi M, Lenzi P, Alessandri MG, Ferrucci M, Fornai F, Paparelli A. Brief and repeated noise exposure produces different morphological and biochemical effects in noradrenaline and adrenaline cells of adrenal medulla. *J Anat*. 2002;200(2):159–68. <https://doi.org/10.1046/j.0021-8782.2001.00014.x>.
33. Yang Y, Zhang E, Zhang J, et al. Relationship between occupational noise exposure and the risk factors of cardiovascular disease in China. *Medicine*. 2018;97(30):e11720. <https://doi.org/10.1097/MD.00000000000011720>.
34. Themann CL, Masterson EA. Occupational noise exposure: a review of its effects, epidemiology, and impact with recommendations for reducing its burden. *J Acoust Soc Am*. 2019;146(5):3879–905. <https://doi.org/10.1121/1.5134465>.
35. Umishio W, Ikaga T, Kario K, et al. Role of housing in blood pressure control: a review of evidence from the Smart Wellness Housing survey in Japan. *Hypertens Res*. 2023;46(1):9–18. <https://doi.org/10.1038/s41440-022-01060-6>.
36. Abu-Saad K, Chetrit A, Eilat-Adar S, et al. Blood pressure level and hypertension awareness and control Differ by Marital Status, Sex, and ethnicity: a Population-based study. *Am J Hypertens*. 2014;27(12):1511–20. <https://doi.org/10.1093/ajh/hpu081>.
37. Lipowicz A, Lopuszanska M. Marital differences in blood pressure and the risk of hypertension among Polish men. *Eur J Epidemiol*. 2005;20(5):421–7. <https://doi.org/10.1007/s10654-005-1752-x>.
38. Joseph NT, Muldoon MF, Manuck SB, et al. The role of Occupational Status in the Association between Job Strain and ambulatory blood pressure during Working and Nonworking days. *Psychosom Med*. 2016;78(8):940–9. <https://doi.org/10.1097/PSY.0000000000000349>.
39. Blumenthal JA, Thyrum ET, Siegel WC. Contribution of job strain, job status and marital status to laboratory and ambulatory blood pressure in patients with mild hypertension. *J Psychosom Res*. 1995;39(2):133–44. [https://doi.org/10.1016/0022-3999\(94\)00087-L](https://doi.org/10.1016/0022-3999(94)00087-L).
40. Karasek R, Baker D, Marxer F, Ahlbom A, Theorell T. Job decision latitude, job demands, and cardiovascular disease: a prospective study of Swedish men. *Am J Public Health*. 1981;71(7):694–705. <https://doi.org/10.2105/AJPH.71.7.694>.
41. Johansson E, Böckerman P, Lundqvist A. Self-reported health versus biomarkers: does unemployment lead to worse health? *Public Health*. 2020;179:127–34. <https://doi.org/10.1016/j.PUHE.2019.10.005>.

42. Essayagh T, Essayagh M, El Rhaffouli A, et al. Prevalence of uncontrolled blood pressure in Meknes, Morocco, and its associated risk factors in 2017. *PLoS ONE*. 2019;14(8):e0220710. <https://doi.org/10.1371/journal.pone.0220710>.

**Publisher's note**

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.